

Issues of Heterotic $(0,2)$ Compactifications

Ralph Blumenhagen

Humboldt Universität Berlin
Institut für Physik, Invalidenstrasse 110, 10115 Berlin, Germany

Abstract: A brief review of some aspects of heterotic $(0,2)$ compactifications in the framework of exactly solvable superconformal field theories and gauged linear sigma models is presented.

1 Introduction

The general class of four dimensional $N = 1$ supersymmetric heterotic compactifications has $(0,2)$ supersymmetry on the world sheet. Geometrically, such models are defined by a stable holomorphic vector bundle (coherent sheaf) over a Calabi-Yau threefold subject to further anomaly cancellation conditions involving the first and second Chern classes of the vector and the tangent bundle. It has long been an open question whether generically such models are indeed consistent vacua of the heterotic string. It has been argued in [1] that $(0,2)$ models realizable as linear sigma models [2] provide a set of exact perturbative vacua. In [3] we presented a class of exactly solvable superconformal field theories with $(0,2)$ supersymmetry, which were argued to describe special points in the moduli space of $(0,2)$ linear sigma models. As a byproduct we got to know a subset of $(0,2)$ models inheriting their defining data from $(2,2)$ models and automatically satisfying all the linear and quadratic anomaly constraints. As a first step towards establishing mirror symmetry in the $(0,2)$ context we defined a way to obtain mirror symmetric pairs at least in the afore mentioned subclass of models [4]. Another approach was made in [5] by successive orbifolding in the Landau-Ginzburg phase.

One is familiar with mirror symmetry, but there exist more general perturbative target space dualities for $(0,2)$ models. Two at large radius completely different looking models can have the same Landau-Ginzburg phase. In [5] it was shown that also at large radius the total dimension of the moduli spaces agrees providing evidence for the conjecture that the models are isomorphic throughout the entire moduli space with a non-trivial map among complex, Kähler and bundle moduli.

Other important approaches to describe $(0,2)$ models using F-theory and equivariant sheaves are not covered here.

2 Exactly solvable SCFTs and Distler-Kachru models

Gepner provided exactly solvable conformal field theories (CFT) describing special point in the moduli space of a Calabi-Yau $(2,2)$ compactification. In this case, up to the application of the bosonic string map the modular invariant partition function is left-right symmetric. In order to find a CFT description of more general $(0,2)$ compactifications

one needs a method for constructing really heterotic partition functions. One way to achieve this is by using simple currents. Since two simple currents can be non-local to each other, the partition function obtained after modding out these simple currents need not to be left-right symmetric. As a generalisation of Gepner models we proposed the following CFTs

	c	$\bar{c}$
flat space-time	2	2
N=2 SCFT	9	9
$(U(1)_2)^{r-3}$	$r-3$	$r-3$
$SO(16-2r) \times E_8$	$16-r$	$16-r$

Table 1: *Ingredients for generalised Gepner models*

In [2] we considered the following modular invariant partition function

$$Z \sim \vec{\chi}(\tau) M(J_{\text{GSO}}) \left(\prod_i M(\Upsilon_i) \right) M(\bar{J}_{\text{GSO}}) \left(\prod_{i=1}^r M(J_i) \right) \left(\prod_{j=1}^{r-3} M(J_{\text{ext}}^j) \right) \vec{\chi}(\bar{\tau}), \quad (1)$$

where the simple currents are chosen in such a way as to guarantee two right moving world sheet supersymmetries, one space time supersymmetry and an extension of the gauge group from $SO(16-2r) \times U(1)^{r-3}$ to E_{9-r} . If the simple current Υ_l does contain factors of both NS and R type, then the left moving supersymmetry is broken and one obtains a model with gauge group $E_{9-r} \times E_8 \times G$. For suitable choices of the simple current, by comparing massless spectra and chiral rings one can identify them as special point in the moduli space of linear sigma models. As an example consider the $(k=3)^5$ Gepner model with $r=4$ and choose

$$\Upsilon = \Phi_{0,-1}^3 \otimes (\Phi_{0,0}^0)^4 \otimes \Phi_{1,2}^{U(1)_2} \otimes \Phi_0^{SO(8)}, \quad (2)$$

having gauge group $SO(10)$ and $N_{16} = 80$ generations, no antigeneration, $N_{10} = 74$ gauge vectors and $N_1 = 350$ gauge singlets. This agrees with the spectrum of the linear sigma model

$$\mathbb{P}_{1,1,1,1,2,2}[4,4] \leftarrow V_{1,1,1,1,1}[5], \quad (3)$$

where the vector bundle V is defined by an exact sequence

$$0 \rightarrow V \rightarrow \bigoplus_{a=1}^5 \mathcal{O}(1) \rightarrow \mathcal{O}(5) \rightarrow 0. \quad (4)$$

Generalising this example in [3] we defined a nice subclass of models. Given a Gepner model with $K_1 = 2\ell - 1$. Let d be the lowest common multiple of the numbers $\{K_i : i = 1, \dots, 5\}$. For models with only four factors set $K_5 = 0$. Then the analysis of the chiral ring reveals that a model obtained by using the following simple currents in the diagonal Gepner parent model

$$\Upsilon = \Phi_{0,-1}^{K_1} \otimes (\Phi_{0,0}^0)^4 \otimes \Phi_{1,2}^{U(1)_2} \otimes \Phi_0^{SO(8)} \quad (5)$$

corresponds to a linear σ -model with the following data

$$\mathbb{P}_{\frac{2d}{2\ell+1}, \frac{\ell d}{2\ell+1}, \frac{d}{K_2+2}, \frac{d}{K_3+2}, \frac{d}{K_4+2}, \frac{d}{K_5+2}} \left[\frac{(\ell+2)d}{2\ell+1}, \frac{2\ell d}{2\ell+1} \right] \leftarrow V_{\frac{d}{2\ell+1}, \frac{d}{K_2+2}, \frac{d}{K_3+2}, \frac{d}{K_4+2}, \frac{d}{K_5+2}} [d]. \quad (6)$$

Roughly speaking one generates $(0, 2)$ data from $(2, 2)$ data automatically satisfying the non-trivial anomaly constraints. This class of models provided a playground for further study. Mirror symmetry has become an important tool in exactly describing moduli spaces of $(2, 2)$ Calabi-Yau compactifications. For $(0, 2)$ models non-perturbative sigma model and target space space corrections are under less control. At least in the class defined above one can generate candidate dual pairs as for instance

$$\mathbb{P}_{1,1,1,1,2,2}[4, 4] \leftarrow V_{1,1,1,1,1}[5] \quad \mathbb{P}_{51,60,80,65,128,128}[256, 256] \leftarrow V_{51,64,60,80,65}[320]. \quad (7)$$

Starting with a $(2, 2)$ mirror pair, one applies the transformation to get two $(0, 2)$ models with still mirror symmetric spectra. Another approach to generate mirror pairs is by orbifolding. To this end we developed orbifold techniques for $(0, 2)$ Landau-Ginzburg models in [4, 5]. We showed that by successive orbifolding of the model in (3) one obtains a mirror symmetric set of models.

3 Target space dualities

Besides mirror symmetry, in the $(0, 2)$ context one can imagine other target space dualities at the perturbative level. It could happen that two models defined by Calabi-Yau threefold and bundle data (M_1, V_1) and (M_2, V_2) are isomorphic as superconformal field theories. One way to realize such a duality exists in the framework of linear sigma models [7]. In the Landau-Ginzburg phase the superpotential reduces to

$$S = \int d^2z d\theta \left[\Gamma^j W_j(\Phi_i) + p \Lambda^a F_a(\Phi_i) \right], \quad \text{with } p = \langle P \rangle, \quad (8)$$

where W_j define the hypersurfaces in a weighed projective space and F_a the bundle. In our former notation this defines a model

$$\mathbb{P}_{\omega_1, \dots, \omega_{N_\omega}}[d_1, \dots, d_{N_d}] \leftarrow V_{n_1, \dots, n_{N_n}}[m]. \quad (9)$$

The parameters ω_i, d_j, n_a, m are related to the $U(1)$ charges of the corresponding superfields $\Phi_i, \Gamma^j, \Lambda^a, P$ in the gauged linear sigma model. In (8) manifold and bundle data appear on equal footing so that it might be possible that two different sets of geometric data lead to the same Landau-Ginzburg models. It was believed for some time that the Landau-Ginzburg point is like a transition point from one $(0, 2)$ model to another [7]. In [6] it was argued that a different scenario occurs, namely that the two models are isomorphic at every point in moduli space. The argument was based on an exact computation of the dimensions of the geometric moduli spaces including complex, Kähler and bundle moduli. As an example consider the quintic

$$\mathbb{P}_4[5] \text{ with deformation of } T \quad (10)$$

and a resolution of

$$\mathbb{P}_{1,1,1,1,1,3}[4, 4] \leftarrow V_{1,1,1,2}[5]. \quad (11)$$

They have the same Landau-Ginzburg locus. Using techniques from toric geometry and homological algebra one can compute the exact dimensions of various cohomology groups. The gauge group in both models is $E_6 \times E_8$. They both have the same number of generations $H^1(M, V) = 101$ and antigerations $H^1(M, V^*) = 1$. For the first model the number of complex, Kähler and bundle moduli is $H^1(M_1, T) = 101$, $H^1(M_1, T^*) = 1$ and $H^1(M_1, \text{End}(T)) = 224$ adding up to a total of 326 moduli. For the second model the

numbers are $H^1(M_1, T) = 86$, $H^1(M_1, T^*) = 2$ and $H^1(M_1, \text{End}(V)) = 238$ amazingly adding up to 326, as well. In all the examples studied, the number of geometric moduli agreed completely where of course the individual contributions of the three kinds of moduli got exchanged. With such high dimensional moduli spaces involved it is difficult to determine the exact map between various moduli. Furthermore, one might ask whether such dualities are of any use for exact non-perturbative computations like for $(2, 2)$ mirror symmetry. All the non-renormalization theorems holding for $(2, 2)$ models are generically not true for $(0, 2)$.

Acknowledgements I would like to thank Andreas Wißkirchen, Rolf Schimmrigk, Sav Sethi and Michael Flohr for their collaboration on part of the work presented in this talk.

References

- [1] E. Witten, *Phases of $N = 2$ theories in two dimensions*, Nucl. Phys. **B403** (1993) 159, hep-th/9301042
J. Distler and S. Kachru, *$(0, 2)$ Landau–Ginzburg theory*, Nucl. Phys. **B413** (1994) 213, hep-th/9309110
- [2] E. Silverstein and E. Witten, *Criteria for conformal invariance of $(0, 2)$ models*, Nucl. Phys. **B444** (1995) 161, hep-th/9503212
- [3] R. Blumenhagen and A. Wißkirchen, *Exactly solvable $(0, 2)$ supersymmetric string vacua with GUT gauge groups*, Nucl. Phys. **B454** (1995) 561, hep-th/9506104
R. Blumenhagen, R. Schimmrigk and A. Wißkirchen, *The $(0, 2)$ exactly solvable structure of chiral rings, Landau–Ginzburg theories and Calabi–Yau manifolds*, Nucl. Phys. **B461** (1996) 460, hep-th/9510055
- [4] R. Blumenhagen, R. Schimmrigk and A. Wißkirchen, *$(0, 2)$ mirror symmetry* Nucl. Phys. **B486** (1997) 598, hep-th/9609167
- [5] R. Blumenhagen and S. Sethi, *On orbifolds of $(0, 2)$ models*, Nucl. Phys. **B491** (1997) 263, hep-th/9611172
R. Blumenhagen and M. Flohr, *Aspects of $(0, 2)$ orbifolds and mirror symmetry*, Phys. Lett. **B404** (1997) 41, hep-th/9702199
- [6] R. Blumenhagen, *Target space duality for $(0, 2)$ compactifications*, Nucl. Phys. **B513** (1998) 573, hep-th/9707198
R. Blumenhagen, *$(0, 2)$ Target Space Duality, CICYs and Reflexive Sheaves* Nucl. Phys. **B514** (1998) 688, hep-th/9710021
- [7] J. Distler and S. Kachru, *Duality of $(0, 2)$ String Vacua*, Nucl. Phys. **B442** (1995) 64, hep-th/9501111
T.-M. Chiang, J. Distler, B. R. Greene, *Some Features of $(0, 2)$ Moduli Space*, Nucl. Phys. **B496** (1997) 590, hep-th/9702030